Egypt


2010 total population: 81 121 077 Income group: Lower middle

NCD mortality		
2008 estimates	males	females
Total NCD deaths (000s)	198.9	172.2
NCD deaths under age 60	38.1	27.8
(percent of all NCD deaths)	00.1	
Age-standardized death rate per 100 000		
All NCDs	829.7	660.0
Cancers	107.3	76.1
Chronic respiratory diseases	33.2	24.3
Cardiovascular diseases and diabetes	427.3	384.0

Behavioural risk factors			
2008 estimated prevalence (%)	males	females	total
Current daily tobacco smoking	35.1	0.5	17.8
Physical inactivity			


Metabolic risk factors			
2008 estimated prevalence (%)	males	females	total
Raised blood pressure	35.5	34.5	35.0
Raised blood glucose	6.2	6.9	6.5
Overweight	60.4	75.3	67.9
Obesity	21.4	44.5	33.1
Raised cholesterol	33.3	43.7	38.6


Proportional mortality (% of total deaths, all ages)





NCDs are estimated to account for 82% of all deaths.

Metabolic risk factor trends









Country capacity to address and respond to NCDs	
Has a Unit / Branch / Dept in MOH with responsibility for NCDs	Yes
There is funding available for: NCD treatment and control NCD prevention and health promotion NCD surveillance, monitoring and evaluation	Yes Yes Yes
National health reporting system includes: NCD cause-specific mortality NCD morbidity NCD risk factors	Yes Yes Yes

No

Has an integrated or topic-specific policy / programme / action plan which is currently operational for:

Cardiovascular diseases	Yes**
Cancer	Yes**
Chronic respiratory diseases	Yes**
Diabetes	Yes**
Alcohol	Yes**
Unhealthy diet / Overweight / Obesity	Yes**
Physical inactivity	Yes**
Tobacco	Yes**

Number of tobacco (m)POWER measures implemented at the highest level of achievement

Has a national, population-based cancer registry

^{... =} no data available

 $^{^{\}star\star}$ = covered by integrated policy/programme/action plan